

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

PCT
WELTORGANISATION FÜR GEISTIGES EIGENTUM
Internationales Büro
INTERNATIONALE ANMELDUNG VERÖFFENTLICHT NACH DEM VERTRAG ÜBER DIE
INTERNATIONALE ZUSAMMENARBEIT AUF DEM GEBIET DES PATENTWESENS (PCT)



<p>(51) Internationale Patentklassifikation ⁵ : A61K 37/02, 47/14, 9/20, 9/48</p>	<p>A1</p>	<p>(11) Internationale Veröffentlichungsnummer: WO 94/23733</p> <p>(43) Internationales Veröffentlichungsdatum: 27. Oktober 1994 (27.10.94)</p>						
<p>(21) Internationales Aktenzeichen: PCT/EP94/01228</p> <p>(22) Internationales Anmeldedatum: 20. April 1994 (20.04.94)</p> <p>(30) Prioritätsdaten:</p> <table style="width: 100%; border: none;"> <tr> <td style="width: 30%;">P 43 12 728.2</td> <td style="width: 30%;">20. April 1993 (20.04.93)</td> <td style="width: 40%;">DE</td> </tr> <tr> <td>P 44 12 201.2</td> <td>8. April 1994 (08.04.94)</td> <td>DE</td> </tr> </table> <p>(71) Anmelder (für alle Bestimmungsstaaten ausser US): DR. RENTSCHLER ARZNEIMITTEL GMBH & CO. [DE/DE]; Mittelstrasse 18, D-88471 Laupheim (DE).</p> <p>(72) Erfinder: und</p> <p>(75) Erfinder/Anmelder (nur für US): FLECK, Monika [DE/DE]; Finkenweg 3, D-88471 Laupheim (DE). NEUER, Klaus [DE/DE]; Gartenstrasse 5, D-88477 Schwendi (DE). WALCH, Hans [DE/DE]; Weißdorn Weg 10, D-88471 Laupheim (DE).</p> <p>(74) Anwälte: SCHWABE, Hans-Georg usw.; Stuntzstrasse 16, D-81677 München (DE).</p>		P 43 12 728.2	20. April 1993 (20.04.93)	DE	P 44 12 201.2	8. April 1994 (08.04.94)	DE	<p>(31) Bestimmungsstaaten: CA, CZ, FI, HU, JP, NO, SK, US, europäisches Patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p>Veröffentlicht <i>Mit internationalem Recherchenbericht. Vor Ablauf der für Änderungen der Ansprüche zugelassenen Frist. Veröffentlichung wird wiederholt falls Änderungen eintreffen.</i></p>
P 43 12 728.2	20. April 1993 (20.04.93)	DE						
P 44 12 201.2	8. April 1994 (08.04.94)	DE						
<p>(54) Title: NEW CYCLOSPORINE-CONTAINING PHARMACEUTICAL COMPOSITIONS FOR ORAL ADMINISTRATION</p> <p>(54) Bezeichnung: NEUE PHARMAZEUTISCHE ZUBEREITUNGEN ZUR ORALEN VERABREICHUNG ENTHALTEND CYCLOSPORIN</p> <p>(57) Abstract</p> <p>New cyclosporine-containing, oral administration pharmaceutical compositions are easy to produce and have a good bioavailability. Besides cyclosporine as active substance, these compositions contain an alkylene polyether or polyester, and may optionally contain an alkylene polyol, an alkylene glycol, a polyalkylene glycol, an alkyl diether or partial ether of a low molecular mono- or polyoxyalkane diol and/or a vegetable oil or its hydrogenated or hydrolyzed product.</p> <p>(57) Zusammenfassung</p> <p>Die Erfindung betrifft neue cyclosporinhalige orale pharmazeutische Zubereitungen. Die neuen pharmazeutischen Zubereitungen sind einfacher herzustellen und verfügen über eine gute Bioverfügbarkeit. Die Zubereitungen enthalten neben dem Wirkstoff Cyclosporin einen Alkylpolyether oder -ester. Fakultativ kann ein Alkylpolyol, ein Alkylglykol, ein Polyalkylglykol, ein Alkyldi- oder Teilether eines niedermolekularen Mono- oder Polyoxyalkandiolis und/oder ein pflanzliches Öl oder dessen hydriertes oder hydrolysiertes Produkt enthalten sein.</p>								

**New Pharmaceutical Preparations Comprising Cyclosporin
for Oral Administration**

The invention relates to new pharmaceutical preparations comprising cyclosporin as active ingredient for oral administration.

Cyclosporins are a class of peptides which are used as immunosuppressants, in particular. Moreover, cyclosporins are known to have antiphlogistic and antiparasitic effects. Therefore, the use of cyclosporins is not limited to immunosuppressants only but relates to all phlogistic diseases including various auto-immune diseases as well as other phlogistic conditions, in particular, phlogistic conditions in which auto-immune processes play a role. The above phlogistic conditions also include, in particular, arthritic diseases such as rheumatoid arthritis as well as rheumatic diseases. Cyclosporins can be used as antiparasitic agents e.g. for the treatment of protozoal infections such as malaria.

Cyclosporins are highly hydrophobic substances, having the consequence that it is difficult to easily process them into pharmaceutical preparations ensuring further sufficient bio-availability. The latter aspect is particularly important, because the cyclosporins possess nephrotoxic side-effects of essential importance. Cyclosporin-containing pharmaceutical preparations proposed so far are based on the use of an alcohol and/or oils or similar vehicles in connection with a surface-active agent. Such preparations are known from DE-OS 29 07 460, for instance. The use of such liquid compositions, however, is accompanied by a number of disadvantages and difficulties. The use of oils or comparable vehicles on oil basis leads to impairment of the sense of taste, in particular, in the case of long-

- 2 -

time administration as a consequence of long-term therapy. Since for dissolving the active ingredient a high amount of alcohol is required, the result will be that in addition the patient is permanently administered alcohol and in the case of evaporation of the alcohol during long-term use the active ingredient precipitates. The attempt to offer such preparations in the form of soft gelatin capsules did not yield a satisfactory solution either due to the higher expenditure connected therewith.

DE-OS 40 03 844 proposes a preparation system which in addition to the active ingredient contains a fatty acid saccharide monoester and a diluent or vehicle by means of which it is said to be possible to provide solid, semi-solid and liquid preparations having a content of cyclosporin in a sufficiently high concentration, so that thus oral administration is comfortably possible and an improved efficiency, for instance, with respect to the bio-availability properties will be achieved. Accordingly, these forms of administration contain at least two components in addition to the active ingredient.

The applicant now has surprisingly found a preparation system for oral administration by means of which it is possible to provide a cyclosporin-comprising pharmaceutical preparation for oral administration, which in addition to the active ingredient cyclosporin contains only one vehicle component. Said component is an alkylene-polyether or alkylene-polyester or any mixture thereof, in which the vehicle system must have an HLB of at least 10. The preparations according to the invention yield a bio-availability of the active ingredient which at least is comparable with the best known cyclosporin-containing preparations.

Having a comparably good bio-availability the pharmaceutical preparations according to the invention can be produced in a more economical way, avoid additives impairing the sense of taste as well as the disadvantageous alcohol contained and, in addition, lead to a better patient compliance within the

- 3 -

sense that the total weight of the formulation to be administered is reduced as compared with known preparations, with the active ingredient concentration staying the same.

Therefore, the invention relates to pharmaceutical preparations for oral administration, containing cyclosporin as active ingredient and being composed as follows:

- a) a cyclosporin as active ingredient,
- b) an alkylene polyether or alkylene-polyester as vehicle or any mixture thereof, with the HLB being at least 10.

Optionally, the preparations according to the invention may contain as further component (c) an alkylene polyole, an alkylene glycole, a polyalkylene glycole, a C₁₂-alkyldiether or partial ether of a lower monooxyalkandiole or polyoxyalkandiole having 2 to 15 carbon atoms and/or a vegetable oil or its hydrated or hydrolysed product.

Moreover, the preparations according to the invention can contain further known, common and pharmaceutically acceptable additives (d) such as are known in the field of the production of oral formulations.

In parts by weight the preparations according to the invention contain 1 to 50 parts by weight of (b) and/or 0.5 to 20 parts by weight of (c) per part by weight active ingredient, preferably 5 to 10 parts by weight (b) and/or 1 to 10 parts by weight (c) per 1 part active ingredient and, in particular 5 parts by weight (b) and/or 1 part by weight (c) per 1 part by weight active ingredient.

In the case of component (b) it suitably pertains to C₁₂ to C₁₈ alkylene-triolether or C₁₂ to C₁₈ alkylene-triester, in particular glycerine. These also include e.g. transesterification products of the alkylene-triolesters with other mono-oles, dioles or polyoles as well as those substances described under "component C³" in DE-OS 40 03 844. Saturated polyglycolised glycerides having an HLB of at least 10 are

- 4 -

particularly advantageous. Preferably, the saturated, polyglycolised glycerides known under the mark term Gelucire (the term Gelucire is a trademark of the company Gattefossé) are used and, in particular, the Gelucire[®] 35/10, 44/14, 42/12, 50/13, 53/10 and any mixtures thereof, in which connection the HLB of the vehicle components used is at least 10.

The optional component (c) comprises, for instance, diethers or partial ethers of lower (C_{2-11}) monooxyalkandioles or polyoxyalkandioles such as are described in DE-OS 39 30 928 in the section relating to the component 1.1. The optional component (c) further comprises C_{1-4} alkylene polyoles, C_{1-4} alkylene glycoles, poly- $(C_{1-4}$ -alkylene)-glycoles, and vegetable oils as well as their hydration and/or hydrolysis products such as castor oil, olive oil, palm oil, coconut oil, corn oil, sesame oil. The component (c) may be contained as single substance or in any mixtures. Preferred examples of the component (c) are glycerine, propylene glycole and polyalkylene glycole having a molecular weight of up to 600, in particular transcitol and castor oil and the hydrated and hydrolysed products thereof.

The further usable additives pertain to pharmaceutically acceptable additives common in the field of oral forms of administration. Examples thereof are the release of controlling substances, thickening agents, preservatives, stabilizers, flavorings, binding agents, lubricants and the like. These additives may amount to up to 50% of the total composition, however, preferably does not exceed 25% and, in particular, not 10% of the total composition.

All of the known natural and synthetic cyclosporins including their analogs and derivatives are suitable for the use in preparations according to the invention. Examples of such cyclosporins are found e.g. in DE-OS 40 03 844 and DE-OS 40 05 190. Preferably cyclosporin A is used.

The oral forms of administration include e.g. liquids, gra-

- 5 -

nulates and solid forms such as tablets and capsules which can be produced according to the common methods known to the person skilled in the art.

The oral forms of administration according to the invention usually are available in standard dose form and contain about 20 to 200 mg, preferably 50 to 100 mg active ingredient per standard dose.

The following examples serve the further illustration of the invention.

Examples

1.	<u>Components</u>	<u>Amount (mg)</u>
	Cyclosporin A	50.0
	Gelucir 53/10	300.0

	Total	350.0
2.	Cyclosporin A	50.0
	Gelucir 44/14	250.0
	Propylene glycole	50.0

	Total	350.0
3.	Cyclosporin A	50.0
	Gelucir 50/13	250.0
	Transcutol	75.0

	Total	375.0
4.	Cyclosporin A	50.0
	Gelucir 44/14	250.0

	Total	300.0

- 6 -

5.	Cyclosporin A	50.0
	Gelucir 50/13	250.0
	Propylene glycole	50.0

	Total	350.0
6.	Cyclosporin A	50.0
	Gelucir 35/10	250.0
	Propylene glycole	25.0

	Total	325.0
7.	Cyclosporin A	50.0
	Gelucir 53/10, 42/12	275.0
	Transcutol	50.0

	Total	375.0
8.	Cyclosporin A	50.0
	Gelucir 42/12	300.0
	Glycerine	25.0

	Total	375.0
9.	Cyclosporin A	50.0
	Gelucir 50/13	250.0
	Castor oil	75.0

	Total	375.0

Production: The compositions of examples 1 to 9 are produced in that the component (b) is melted by heating preferably to at least 60°C and the active ingredient (a) is dissolved therein by stirring. If desired, optional component (c) is added to the melted mass.

- 7 -

Subsequently, the preparations obtained are filled, for instance, in liquid form into hard-gelatin capsules of the desired size in the concentrations desired. The compositions can also be further processed to tablets in the known manner. For this purpose, the melted masses are produced as described in the above. The liquid melted masses are poured out and after solidification diminished by means of a sieving machine. The granulates produced such are mixed with the usual adjuvants such as slip agents and lubricants, blasting agents, fillers, flavor corrigents etc. The finished mixtures are pressed to tablets having the desired content of cyclosporin. The tablets may also be coated with a protective cover.

Bio-Availability:

Examinations as to the bio-availability of the compositions according to the invention on dogs.

A group of six Beagle dogs was used for the bio-availability examinations. The test drugs were orally applied to the animal with an empty stomach by means of oesophageal sounds. At defined times blood is taken from the Vena saphena of the animals and collected in corresponding plastic tubes with EDTA additive. The blood samples are stored until assaying at -18°C . Assay of the cyclosporin takes place in the whole blood by means of fluorescence-polarisation immunoassay (FPIA).

The areas under the curves (AUC) in which the blood drug concentration is applied relative to time were calculated according to the trapezoid rule. The average AUC values of compositions according to the invention are shown in the following table in comparison to the commercially available substances of cyclosporin drinking solution and cyclosporin capsules (Sandimmun[®]) which were ascertained in the same manner in the same dosage with the same dogs.

- 8 -

<u>Examples</u>	<u>AUC (0-12 h) ng/ml</u>
Example 2	9035 + 2134
Example 3	7859 + 1512
Example 4	7552 + 1194
Example 5	8228 + 857
Cyclosporin Drinking Solution	7980 + 1320
Cyclosporin Capsules	8098 + 1504

As the above tests on the bio-availability show, the pharmaceutical preparations according to the invention make it possible to provide the active ingredient cyclosporin in such an oral form that its bio-availability at least corresponds to the preparations known so far.

Patent Claims

1. Pharmaceutical preparation for oral administration containing as the only component or consisting of
 - (a) a cyclosporin as active ingredient, and
 - (b) an alkylene-polyether or alkylene-polyester either alone or in any mixture as vehicle, whereby the HLB of the component (b) used being at least 10.
2. Pharmaceutical preparation according to claim 1, further containing (c) an alkylene-polyole, alkylene-glycole, a polyalkylene-glycole, an alkyl-diether or partial ether of a lower monooxyalkandiol or polyoxyalkandiol and/or a vegetable oil or its hydrated or hydrolysed product either alone or in any mixtures.
3. Preparation according to claim 1 or 2, in which the respective components (a), (b) and/or (c) are available in the following weight ratios: 1:1-50:0.5-20, preferably 1:5-10:1-10, in particular 1:5:1.
4. Preparation according to one of claims 1 to 3, in which the component (b) is chosen from among saturated polyglycolised glycerides.
5. Pharmaceutical preparation according to the previous claim 4, in which the component (b) is chosen from among the Gelucires Gelucir[®] 35/10, 44/14, 42/12, 50/13, 53/10 and any mixtures thereof.
6. Pharmaceutical preparation according to claim 2, in which the additional component (c) is chosen from among glycerine, propylene glycole, PEG with MG up to approx. 600,

- 10 -

transcutol and castor oil.

7. Pharmaceutical preparation according to one of the previous claims in the form of hard-gelatin capsules or in the form of a tablet.

8. Pharmaceutical preparation according to one of the previous claims, characterized in that the active ingredient concentration is 20 to 200 mg, preferably 50 to 100 mg per dose unit.

- 11 -

Summary

The invention relates to new cyclosporin-comprising oral pharmaceutical preparations. The new pharmaceutical preparations can be produced more easily and have a good bio-availability. In addition to cyclosporin as active ingredient the preparations contain an alkylene-polyether or alkylene-polyester. Optionally, an alkylene-polyole, an alkylene-glycole, a polyalkylene-glycole, an alkyldiether or partial ether of a lower monooxyalkandiole or polyoxyalkandiole and/or a vegetable oil or its hydrated or hydrolysed product may be contained.

**New Pharmaceutical Preparations Comprising
Cyclosporin for Oral Administration**

BACKGROUND OF THE INVENTION

1. Field of the Invention

The invention relates to new pharmaceutical preparations comprising cyclosporin as active ingredient for oral administration.

2. Description of the Prior Art

Cyclosporins are a class of peptides which are used as immunosuppressants, in particular. Moreover, cyclosporins are known to have antiphlogistic and antiparasitic effects. Therefore, the use of cyclosporins is not limited to immunosuppressants only but relates to all phlogistic diseases including various auto-immune diseases as well as other phlogistic conditions, in particular, phlogistic conditions in which auto-immune processes play a role. The above phlogistic conditions also include, in particular, arthritic diseases such as rheumatoid arthritis as well as rheumatic diseases.

Cyclosporins can be used as antiparasitic agents e.g. for the treatment of protozoal infections such as malaria.

Cyclosporins are highly hydrophobic substances, having the consequence that it is difficult to easily process them into pharmaceutical preparations ensuring further sufficient bio-availability. The latter aspect is particularly important, because the cyclosporins possess

- 2 -

nephrotoxic side-effects of essential importance. Cyclosporin-containing pharmaceutical preparations proposed so far are based on the use of an alcohol and/or oils or similar vehicles in connection with a surface-active agent. Such preparations are known from DE-OS 29 07 460, for instance. The use of such liquid compositions, however, is accompanied by a number of disadvantages and difficulties. The use of oils or comparable vehicles on oil basis leads to impairment of the sense of taste, in particular, in the case of long-time administration as a consequence of long-term therapy. Since for dissolving the active ingredient a high amount of alcohol is required, the result will be that in addition the patient is permanently administered alcohol and in the case of evaporation of the alcohol during long-term use the active ingredient precipitates. The attempt to offer such preparations in the form of soft gelatin capsules did not yield a satisfactory solution either due to the higher expenditure connected therewith.

DE-OS 40 03 844 proposes a preparation system which in addition to the active ingredient contains a fatty acid saccharide monoester and a diluent or vehicle by means of which it is said to be possible to provide solid, semi-solid and liquid preparations having a content of cyclosporin in a sufficiently high concentration, so that thus oral administration is comfortably possible and an improved efficiency, for instance, with respect to the bio-availability properties will be achieved. Accordingly, these forms of administration contain at least two components in addition to the active ingredient.

SUMMARY OF THE INVENTION

The applicant now has surprisingly found a preparation system for oral administration by means of which it is possible to provide a cyclosporin-comprising pharmaceutical prepara-

- 3 -

tion for oral administration, which in addition to the active ingredient cyclosporin contains only one vehicle component. Said component is an alkylene-polyether or alkylene-polyester or any mixture thereof, in which the vehicle system must have an HLB of at least 10. The preparations according to the invention yield a bio-availability of the active ingredient which at least is comparable with the best known cyclosporin-containing preparations.

Having a comparably good bio-availability the pharmaceutical preparations according to the invention can be produced in a more economical way, avoid additives impairing the sense of taste as well as the disadvantageous alcohol contained and, in addition, lead to a better patient compliance within the sense that the total weight of the formulation to be administered is reduced as compared with known preparations, with the active ingredient concentration staying the same.

Therefore, the invention relates to pharmaceutical preparations for oral administration, containing cyclosporin as active ingredient and being composed as follows:

- a) a cyclosporin as active ingredient,
- b) an alkylene polyether or alkylene-polyester as vehicle or any mixture thereof, with the HLB being at least 10.

Optionally, the preparations according to the invention may contain as further component (c) an alkylene polyole, an alkylene glycole, a polyalkylene glycole, a C_{2-15} -alkyldiether or partial ether of a lower monooxyalkandiole or polyoxyalkandiole having 2 to 15 carbon atoms and/or a vegetable oil or its hydrated or hydrolysed product.

- 4 -

Moreover, the preparations according to the invention can contain further known, common and pharmaceutically acceptable additives (d) such as are known in the field of the production of oral formulations.

In parts by weight the preparations according to the invention contain 1 to 50 parts by weight of (b) and/or 0.5 to 20 parts by weight of (c) per part by weight active ingredient, preferably 5 to 10 parts by weight (b) and/or 1 to 10 parts by weight (c) per 1 part active ingredient and, in particular 5 parts by weight (b) and/or 1 part by weight (c) per 1 part by weight active ingredient.

In the case of component (b) it suitably pertains to C_3 to C_6 alkylene-triolether or C_3 to C_6 alkylene-triolester, in particular glycerine. These also include e.g. transesterification products of the alkylene-triolesters with other mono-oles, dioles or polyoles as well as those substances described under "component C³" in DE-OS 40 03 844. Saturated polyglycolised glycerides having an HLB of at least 10 are particularly advantageous. Preferably, the saturated, polyglycolised glycerides known under the mark term Gelucire (the term Gelucire is a trademark of the company Gattefossé) are used and, in particular, the Gelucire[®] 35/10, 44/14, 42/12, 50/13, 53/10 and any mixtures thereof, in which connection the HLB of the vehicle components used is at least 10.

The optional component (c) comprises, for instance, diethers or partial ethers of lower (C_{2-12}) monooxyalkandioles or polyoxyalkandioles such as are described in DE-OS 39 30 928 in the section relating to the component 1.1. The optional component (c) further comprises C_{2-6} alkylene polyoles, C_{2-6} alkylene glycoles, poly- $(C_{2-6}$ -alkylene)-

- 5 -

glycoles, and vegetable oils as well as their hydration and/or hydrolysis products such as castor oil, olive oil, palm oil, coconut oil, corn oil, sesame oil. The component (c) may be contained as single substance or in any mixtures. Preferred examples of the component (c) are glycerine, propylene glycole and polyalkylene glycole having a molecular weight of up to 600, in particular transcutool and castor oil and the hydrated and hydrolysed products thereof.

The further usable additives pertain to pharmaceutically acceptable additives common in the field of oral forms of administration. Examples thereof are the release of controlling substances, thickening agents, preservatives, stabilizers, flavorings, binding agents, lubricants and the like. These additives may amount to up to 50% of the total composition, however, preferably does not exceed 25% and, in particular, not 10% of the total composition.

All of the known natural and synthetic cyclosporins including their analogs and derivatives are suitable for the use in preparations according to the invention. Examples of such cyclosporins are found e.g. in DE-OS 40 03 844 and DE-OS 40 05 190. Preferably cyclosporin A is used.

The oral forms of administration include e.g. liquids, granulates and solid forms such as tablets and capsules which can be produced according to the common methods known to the person skilled in the art.

The oral forms of administration according to the invention usually are available in standard dose form and contain about 20 to 200 mg, preferably 50 to 100 mg active ingredient per standard dose.

- 6 -

DESCRIPTION OF THE PREFERRED EMBODIMENTS

The following examples serve the further illustration of the invention.

Examples

1.	<u>Components</u>	<u>Amount (mg)</u>
	Cyclosporin A	50.0
	Gelucir 53/10	300.0

	Total	350.0
2.	Cyclosporin A	50.0
	Gelucir 44/14	250.0
	Propylene glycole	50.0

	Total	350.0
3.	Cyclosporin A	50.0
	Gelucir 50/13	250.0
	Transcutol	75.0

	Total	375.0
4.	Cyclosporin A	50.0
	Gelucir 44/14	250.0

	Total	300.0

- 7 -

5.	Cyclosporin A	50.0
	Gelucir 50/13	250.0
	Propylene glycole	50.0

	Total	350.0
6.	Cyclosporin A	50.0
	Gelucir 35/10	250.0
	Propylene glycole	25.0

	Total	325.0
7.	Cyclosporin A	50.0
	Gelucir 53/10, 42/12	275.0
	Transcutol	50.0

	Total	375.0
8.	Cyclosporin A	50.0
	Gelucir 42/12	300.0
	Glycerine	25.0

	Total	375.0
9.	Cyclosporin A	50.0
	Gelucir 50/13	250.0
	Castor oil	75.0

	Total	375.0

Production: The compositions of examples 1 to 9 are produced in that the component (b) is melted by heating preferably to at least 60°C and the active ingredient (a) is dissolved

- 8 -

therein by stirring. If desired, optional component (c) is added to the melted mass.

Subsequently, the preparations obtained are filled, for instance, in liquid form into hard-gelatin capsules of the desired size in the concentrations desired. The compositions can also be further processed to tablets in the known manner. For this purpose, the melted masses are produced as described in the above. The liquid melted masses are poured out and after solidification diminished by means of a sieving machine. The granulates produced such are mixed with the usual adjuvants such as slip agents and lubricants, blasting agents, fillers, flavor corrigents etc. The finished mixtures are pressed to tablets having the desired content of cyclosporin. The tablets may also be coated with a protective cover.

Bio-Availability:

Examinations as to the bio-availability of the compositions according to the invention on dogs.

A group of six Beagle dogs was used for the bio-availability examinations. The test drugs were orally applied to the animal with an empty stomach by means of oesophageal sounds. At defined times blood is taken from the Vena saphena of the animals and collected in corresponding plastic tubes with EDTA additive. The blood samples are stored until assaying at -18°C . Assay of the cyclosporin takes place in the whole blood by means of fluorescence-polarisation immunoassay (FPIA).

The areas under the curves (AUC) in which the blood drug concentration is applied relative to time were calculated

- 9 -

according to the trapezoid rule. The average AUC values of compositions according to the invention are shown in the following table in comparison to the commercially available substances of cyclosporin drinking solution and cyclosporin capsules (Sandimmun®) which were ascertained in the same manner in the same dosage with the same dogs.

<u>Examples</u>	<u>AUC (0-12 h) ng/ml</u>
Example 2	9035 + 2134
Example 3	7859 + 1512
Example 4	7552 + 1194
Example 5	8228 + 857
Cyclosporin Drinking Solution	7980 + 1320
Cyclosporin Capsules	8098 + 1504

As the above tests on the bio-availability show, the pharmaceutical preparations according to the invention make it possible to provide the active ingredient cyclosporin in such an oral form that its bio-availability at least corresponds to the preparations known so far.

- 10 -

Patent Claims

1. Pharmaceutical preparation for oral administration containing as the only component or consisting of
 - (a) a cyclosporin as active ingredient, and
 - (b) an alkylene-polyether or alkylene-polyester either alone or in any mixture as vehicle, whereby the HLB of the component (b) used being at least 10.
2. Pharmaceutical preparation according to claim 1, further containing (c) an alkylene-polyole, alkylene-glycole, a polyalkylene-glycole, an alkyl-diether or partial ether of a lower monooxyalkandiol or polyoxyalkandiol and/or a vegetable oil or its hydrated or hydrolysed product either alone or in any mixtures.
3. Preparation according to claim 1, in which the respective components (a), (b) and/or (c) are available in the following weight ratios: 1:1-50:0.5-20, preferably 1:5-10:1-10, in particular 1:5:1.
4. Preparation according to claim 1, in which the component (b) is chosen from among saturated polyglycolised glycerides.
5. Pharmaceutical preparation according to the previous claim 4, in which the component (b) is chosen from among the Gelucires Gelucir[®] 35/10, 44/14, 42/12, 50/13, 53/10 and any

- 11 -

mixtures thereof.

6. Pharmaceutical preparation according to claim 2, in which the additional component (c) is chosen from among glycerine, propylene glycole, PEG with MG up to approx. 600, transcitol and castor oil.

7. Pharmaceutical preparation according to claim 1 in the form of hard-gelatin capsules or in the form of a tablet.

8. Pharmaceutical preparation according to claim 1, in which the active ingredient concentration is 20 to 200 mg, preferably 50 to 100 mg per dose unit.

- 12 -

Summary

The invention relates to new cyclosporin-comprising oral pharmaceutical preparations. The new pharmaceutical preparations can be produced more easily and have a good bio-availability. In addition to cyclosporin as active ingredient the preparations contain an alkylene-polyether or alkylene-polyester. Optionally, an alkylene-polyole, an alkylene-glycole, a polyalkylene-glycole, an alkyl-diether or partial ether of a lower monooxyalkandiol or polyoxyalkandiol and/or a vegetable oil or its hydrated or hydrolysed product may be contained.